

Insect pest diversity of standing crops and traditional pest management in agricultural areas of Mandakini Valley, Garhwal Himalaya, Uttarakhand, India

Vandana Mehrwar*, V. P. Uniyal

Department of Landscape Level Planning and Management, Wildlife Institute of India, Chandrabani, Dehradun-248001, Uttarakhand, India

Received: 03 Jun 2020; Received in revised form: 02 Jul 2021; Accepted: 10 Jul 2021; Available online: 18 Jul 2021

©2021 The Author(s). Published by AI Publications. This is an open access article under the CC BY license

(<https://creativecommons.org/licenses/by/4.0/>)

Abstract— The mountain farming communities of the Garhwal Himalaya rely on a conventional approach to agriculture to meet their subsistence needs. The resilience of local crop varieties plays a significant role in crop productivity in the indigenous agricultural system. In such circumstances, the protection of the crop from insect pests becomes paramount. Traditional ecological knowledge plays a crucial role in safeguarding standing crops from production losses in an environmentally benign and sustainable manner. The investigators in this study have surveyed the Mandakini valley to document the indigenous practices undertaken by the farming folks to protect the crops from pest infestation in the region. These practices are discoursed here and, further, look into the potential of natural predators as bio-control. The findings indicated that pests from the order Coleoptera had the most species, followed by Lepidoptera and Hemiptera. Most of the pests in the study site were serious defoliators, damaging the young foliage of the crops. Some entirely fed upon their roots-stems, while the rest were leaf miners and sapsuckers, thus compromising the overall well-being of the plant. In a developing country like India, there is a lack of reliable data that sheds light on the annual crop losses incurred by these pests. Thus, it becomes pertinent to compute an overall estimate of crop losses at various stages of crop production, from seed storage to post-harvest times.

Keywords— Crop loss, Insect pest, Mandakini Valley, Natural predators, Pest management.

I. INTRODUCTION

FAO/WHO (2014) has defined a pest as "any species, strain or biological type of plant, creature or pathogenic agent that damages plants or parts of them and incorporates vectors of parasites or pathogens of human and animal infections and creatures, causing a public health nuisance." Every year on June 6th, "World Pest Awareness Day" is observed to raise awareness about how pest management contributes to human well-being and their survival. Insects are the most ubiquitous, diverse, and abundant animal group on the planet. These small, versatile beings are the major contenders for food and other useful resources for humans that are produced in the course of farming (Oerke & Dehne 2004). The Food and Agriculture Organization (FAO) has labelled 2020 as the "International Year of Plant Health," with the purpose of raising awareness about how plant health may aid in hunger

prevention, poverty alleviation, environmental protection, and economic development. Plants constitute at least 80 percent of our nourishment, yet they are constantly threatened by pests and diseases (FAO 2019). Pests and diseases destroy up to 40% of the world's food crops each year. This incurs an annual agricultural trade loss of more than \$220 billion, and results in hunger, and eventually interferes with rural income. It is reckoned that globally, food crops are harmed by over 10,000 insect species, 30,000 weed species, 100,000 diseases, and 1,000 nematode species (Dhaliwal et al. 2007).

The first systematic effort to assess crop losses due to different pests at a global level was accomplished by Cramer (1967), who projected total annual losses to be around 34% for major crops and vegetables. It was also specified explicitly that one-third of the total crop production worldwide is spoiled due to weeds, insects, and

diseases. (Oerke 2006). Losses due to insect pests in the context of Indian agriculture have also been reviewed in the past (Pradhan 1964; Krishnamurthy Rao & Murthy 1983; Atwal 1986; Jayaraj 1993; Lal 1996; Dhaliwal & Arora 1996, 2002; Dhaliwal et al. 2003, 2004), and crop losses after the green revolution era were relatively higher than those recorded globally (Pradhan 1964; Dhaliwal et al. 2004). Crop losses increased from 7.2 percent in the early 1960s to 23.3 percent in the early 2000s, but then fell to 17.5 percent in the 21st (Dhaliwal et al. 2007; Dhaliwal et al. 2010). Agriculture is the mainstay of the people residing in the rural areas of the Mandakini valley. Agriculture or allied practices employ more than 75% of the total population, which is substantially dominated by the subsistence mode of farming. The conventional method of mountain agriculture is the only way to meet the subsistence needs of the farming communities in the Garhwal Himalaya. Food grains like paddy, wheat, millet, barley, oil seeds, lentils, and vegetables such as *Solanum tuberosum*, *Solanum lycopersicum*, *Allium cepa*, *Allium sativum*, *Capsicum annum*, *Solanum melongena*, *Brassica oleracea*, *Cucurbits*, *Raphanus sativus*, *Pisum sativum*, *Zingiber officinale* etc., are widely grown in the region. Smallholders and marginal farmers feed more than 2/3rds of the population, but pests destroy at least 30%-40% of the food crops they grow. However, the resilience of local crop varieties plays a significant role in overall farm productivity and in such circumstances, crop protection from insect pests becomes paramount. Traditional ecological knowledge plays a crucial role in safeguarding standing crops from production losses in an environmentally benign and sustainable manner.

This investigation comprising insect pests of standing crops is a pioneer in the region. Earlier studies involving insect pests of stored grains were conducted and evaluated by research investigators in Rudraprayag district. More than 50% of insect mortality for stored wheat grains was demonstrated using smoke generated from neem leaves and cow dung burning, which is both cost-effective and eco-friendly (Yadav & Tiwari 2018).

Cutworms, root weevils, moth and beetle larvae, fruit flies, fruit or shoot miners, plant hoppers, and mealy bugs are major defoliators, pod damagers, sap suckers, shoot borers, and root feeders that destroy crops from seedling to maturity (Plate 1). The mechanism for averting the pest population explosion is crop diversification in the agro-ecosystem (Rao et al. 2015). The motive behind the study is to get a quick overview of insects as major crop pests, examine their extent of damage to agricultural crops, review existing control measures, and describe potential natural predators as bio-control in the region. Scholars, scientists, researchers, and policymakers will benefit from

the study since it sets the path for further investigation and development of a sustainable approach to protecting crops in mountain agro-biodiversity.

II. STUDY AREA

The research was conducted in the villages of the Mandakini valley, in the Rudraprayag district of the Garhwal region. These villages appear to occur in clusters or hamlets and lie between 1654-2029 metres, surrounded by forests. The detailed account of their geographical coordinates, area, population, and number of households as per census 2011 is specified below (Fig. 1).




Fig 1: Study sites of the Mandakini valley

Broadly, the seasons are divided into three, viz., summer (April-June), the monsoon (July-September), and winter (November-March). Summers are pleasantly mild while winters are generally cold and prolonged with snowfall. Rains are mostly confined to the rainy season and heavy downpours in the rainy season frequently cause landslides and soil erosion. The maximum monthly temperature in the area varies from around 19° C to 28° C. The vegetation is a temperate broad leaf type (moist deciduous/ evergreen/ mixed) with dominant species such as *Quercus*, *Rhododendron*, *Pinus*, *Aesculus*, *Acer*, *Juglans*, *Thamnocalamus*, *Daphniphyllum*, *Prunus*, *Myrica*, and others. Despite feasible climatic conditions, coarse, and well-drained acidic soil, the net yield is low. Farmers are unable to meet their food needs due to erratic weather, a lack of irrigation facilities, mono-cropping, non-laboratory soil testing, insect pest attacks, and crop raiding by wild animals. As a result, the majority of people rely on local

stores and the Public Distribution System (PDS) to meet their daily calorie needs.

III. METHODOLOGY

Three years of research was conducted on the farms of three Gram Panchayats in the Mandakini valley, which included seven villages. Periodical monitoring of insects pests of farmland, homesteads, kitchen gardens, and poly houses was carried out during the pre and post-monsoon seasons from 2017 to 2019. In this study, a random household survey was conducted in each village using a semi-structured questionnaire set, key respondents, and keen observation to enumerate the area under crop cultivation, crop composition, cropping pattern, crop pests, and diseases of cultivated crops. Informal dialogues with knowledgeable family members, particularly women, who are actively involved in agricultural activities, were used to gather the information.

A few community-based discussions were also held, mostly about recent farming trends, insecticide/pesticide use, traditional knowledge, and future aspects of farming related to climate change. Sampling methods such as opportunistic sampling and aerial sampling via sweep netting, handpicking, and ground digging were done as per the study needs. Specimens were documented either through photography or sample collection, and dry pinned for further identification and research. The information acquired from natives was analyzed in order to investigate scientific rationality.

IV. RESULT

A total of 187 respondents were interviewed and data was recorded from each village in the Ukhimath block. Obtained data was compiled and analyzed (Fig. 2 & Table 1). The residents of the Mandakini Valley were well aware of the insect pests present on their farmland, but they generally used traditional methods as control measures. Farmers used trap crops, cover crops, farmyard manure, and a mixture of salt-burnt fuel wood chullah ashes to thwart the advent of insect pest infestations on the standing crops.

Beetles are primarily crop pests of cereals, oilseeds, fruits, vegetables, and stored grains (Patole 2017). The result exhibited major farm insects as pests, comprising 32 genera from 22 different families from the study area. Findings indicated that pests from the order Coleoptera (14 sp.) were the maximum in number, followed by Lepidoptera (07 species), Hemiptera (05 sp.), and Diptera (03 sp.) while Orthoptera, Thysanoptera, and Dermaptera contributed with single species respectively.

Coleoptera (Beetles) were dominated by Chrysomelidae (05 sp.), Scarabaeidae (03), and Elateridae (02), while Nitidulidae, Meloidae, Brentoidae, and Curculionidae families each contributed a single species. Hemiptera (True bugs) have six families, including Coreidae (02 sp), Aphidae, Miridae, Pseudococcidae, Flatidae, and Pentatomidae, each with one sp. Lepidoptera (Butterflies and Moths) comes in second with four families of Noctuidae (04 species), while Erebidae, Sphingidae, and Papilionidae each have one species; Diptera has three families of Tipulidae, Psychodidae, and Tephritidae (one sp. each). A bar graph chart depicts a detailed account of these orders and families, along with the number of species (Fig. 2).




Fig.2: Pie chart showing the families belonging to various orders.

Larvae of the orders Lepidoptera, Coleoptera, and Diptera, are voracious feeders and are considered phytophagous pests. Braconids and Ichneumonidae are internal or external parasitoids of these pests that help in checking their population naturally (Plate 2). Birds, rodents, beetles, and earwigs are common predators of larval and adult insect pests. Beetles (larval stage) and weevils are generally root feeders and often damage the flowers and foliage while reaching for pollen and nectar. Most pests are serious defoliators, damaging the young foliage of the crops. Some entirely fed upon their roots-stems, while the rest were leaf miners and sapsuckers, thus jeopardising the overall well-being of the plant. *Altica himalayensis*, *Cotinis nitida*, *Phyllophaga* sp, *Holotrichia* sp, *Bactrocera cucurbitae*, *Aphis* sp, *Pieris brassicae*, *Clavicornis* sp., *Trichoplusia ni*, *Agrotis ipsilon*, and *Thrips tabaci*, are the most common pests that severely affect standing crops and vegetables. The detailed accounts of the major pests of crops about their classification, host plants, and nature of damage are presented below (Table 1 & Plate 1).

V. DISCUSSION

Indigenous farming systems use Traditional Ecological Knowledge (TEK) to control insect pest infestations, such as deep ploughing to expose eggs and larvae of pests, spreading completely decomposed Farm Yard Manure (FYM) on prepared land, dusting of salt-burnt fuelwood chullah ash, mixture of cow dung-urine on standing crops, and stubble burning to clear leftover residues after harvesting to get rid of the exposed pests beneath the soil (Chandola et al. 2011). Leaf litter and livestock feed collected from forests are the primary sources of traditional FYM in the hill regions of Uttarakhand (Maikhuri et al. 2015). In the long run, narrow/selective spectrum chemical pesticides are preferred over broad-spectrum pesticides.

78.5 % of flowering plants in temperate habitats require an animal pollinator to successfully reproduce. Wild entomofauna and birds largely contribute to the productivity of crops through the provision of ecosystem services, such as pollination and natural pest control (Classen et al. 2014). There have been numerous reports of parasitic and predatory natural enemies being used to control agricultural insect pests (Van den Bosch et al. 1982). Spiders are regarded as important predators that aid in the regulation of insect pest population densities (Pickett et al. 1946; Dondale 1956; Kajak et al. 1968; Fox & Dondale 1972; Tanaka 1989). Biological control involves natural predators and parasitoids that are represented by the entomophagus groups (Sampaio et al. 2009). It is often used as a management tool in Integrated Pest Management (IPM), which is economically feasible and has a minimal environmental impact. Many insect pests have their natural enemies, like Braconids (Parasitoid- cutworm wasp), Micromus (Brown lacewing), Coccinellids (Ladybird beetles), Syrphidae (Hover flies), Soldier beetles, Spiders, and Capoletis (Parasitoid wasp-Ichneumonidae) (Plate 2).

Flea beetles can be deterred by several traps and companion plants (such as Basils, Beans, Brassica sp., Secale cereal, Zea mays, Cucurbits, Tagetes sp., Trifolium sp., wild Alliums, Mentha sp., Juglans sp.) that can be intercropped with the primary crop or planted at the periphery of cultivated land. The majority of insects provide direct or indirect benefits to humans (Peters 1993), either in the form of pollinators, insect predators or parasites of pests. The wildflower strip elevates farmland biodiversity, enhances foraging opportunities for various insect pests and pollinators, and also ensures improved productivity (Matthias et al. 2016). Thus, pest control and pollination services are complementary in nature and essential for the sustainability of the mountain agroecosystem.

Recently, the locust invasion of the Rabi crops in Pakistan and the Middle East led to a total crop failure, while in Africa, the situation was so grave that Somalia declared a national emergency. A similar onslaught occurred in Rajasthan, Gujarat, Maharashtra, Madhya Pradesh, and western Uttar Pradesh, galloping several lakh hectares of Rabi crop. Every year, an unprecedented amount of stored food grains, vegetables, and standing crops are destroyed by the infliction of these pests. In a developing country like India, there is a lack of reliable data that sheds light on the annual crop losses incurred by these pests. Thus, it becomes pertinent to compute an overall estimate of crop losses at various stages of crop production, from seed storage to post-harvest times. Emphasis should be given to different eco-friendly methods, such as the use of natural enemies, predators, cultural practices, bio-control, insect-resistant varieties, and the use of scientifically proven transgenic crops (Plate 2). The dissemination of farmer-friendly information about pest management via different platforms, such as Kisan web portal, KVKs (Krishi Vigyan Kendras), Krishi melas, social media, newsletters, pamphlets, and so on, ensures rapid adoption by rural folks and averts crop damage. Therefore, it is imperative to prioritize our food security program to contend with our limited resources and rising population.

VI. CONCLUSION

A total of 35 insect pest individuals from seven orders with a total of 23 families were documented from the study area. Pests from the order Coleoptera (14 species) were found to be the highest in number, followed by Lepidoptera and Hemiptera (07 sp. each), and Diptera (03 sp.). Coleoptera (07), with the maximum number of families, surpassed the others, followed by Hemiptera (06), Lepidoptera (04), and Diptera (03). Chrysomelidae dominated in terms of pest species number, followed by Noctuidae, Scarabaeidae, Acrididae, Elateridae, Coreidae, Brentidae, Curculionidae, Meloidae, Nitidulidae, Forficulidae, Psychodidae, Tephritidae, Tipulidae, and Aphididae. Globally, only 1% of all insects are pests (Triplehorn & Johnson 2005), but they are responsible for the loss of 13% of crop productivity and 9% of forest production (Pimental et al. 2000). Moths and butterflies are beneficial as pollinators, but their larvae are potentially harmful. Cabbage White butterfly larvae, an invasive species, are serious pests of Brassicaceae plants (Snell-Rood & Papaj 2009, Cipollini 2002). The majority of the insect pests of standing crops were Coleopteran beetles, larvae of Lepidopterans and Hemipterans, of which *Altica himalayensis*, *Cotinis nitida*, *Phyllophaga sp.*, *Holotrichia sp.*, *Bactrocera cucurbitae*,

Aphis sp., *Clavicornis* sp., *Trichoplusia ni*, *Agrotis* sp., and *Thrips* damaged the crops severely.

ACKNOWLEDGMENTS

We sincerely acknowledge the Director and Dean, Wildlife Institute of India (WII) for their encouragement and guidance. We are grateful to the local people of the study villages for sharing their valuable knowledge and help in data collection. We would like to put down our sincere thanks to the Ministry of Environment, Forest & Climate Change (MoEF & CC), Govt. of India and G.B. Pant National Institute of Himalayan Environment & Sustainable Development (GBPNIHESD), Almora for providing financial assistance in undertaking this work.

REFERENCES

- [1] D. Pickett, N. A. Patterson, H. T. Stultz, and F. T. Lord, "The influence of spray programs on the fauna of apple orchards in Nova Scotia: An appraisal of the problem & a method of approach," *Scient. Agric.*, 1946, 26: 590-600.
- [2] A. Classen, P. Marcell, S. W. Ferger, M. H. Bonitz, J. M. Schmack, G. Maassen, M. Schleuning, E. K. V. Kalko, K. B. Gaese, and I. S. Dewenter, "Complementary ecosystem services provided by pest predators and pollinators increase quantity and quality of coffee yields," *Proc. R. Soc. B.*, 2014, 281:20133148. <http://dx.doi.org/10.1098/rspb.2013.3148>. Accessed 19 January 2021
- [3] A. Kajak, L. Andrzejewska, and Z. Wojcik, "The role of spiders in the decrease of damages caused by Acridoidea on meadows - Experimental investigations," *Ekol. pol.*, 1968, 16: 755-764.
- [4] A. S. Atwal, "Future of pesticides in plant protection," *Proc Indian Natn Sci Acad.*, 1986, 52:77-90.
- [5] C. A. Triplehorn, and N. F. Johnson, "*Borror and DeLong's Introduction to the Study of Insects*," 7th Edition. Belmont, CA: Brooks/Cole, Thomson Learning, 2005.
- [6] C. D. Dondale, "Annotated list of spiders (Araneae) in Nova Scotia apple orchards," *Can. Entom.*, 1956, 88: 697-700.
- [7] C. J. S. Fox and C. D. Dondale, "Annotated list of spiders (Araneae) from hayfields and their margins in Nova Scotia," *Canadian Entomologist*, 1972, 104: 1911-1915.
- [8] D. Cipollini, "Variation in the expression of chemical defenses in *Alliaria petiolata* (Brassicaceae) in the field and common garden," *American Journal of Botany*, 2002, 89, 1422-1430.
- [9] D. Pimental, "Environmental and economic costs of non-indigenous species in the United States," *BioScience*, 2000, 50, 53-65.
- [10] E. C. Oerke and H. W. Dehne, "Safeguarding production – losses in major crops and the role of crop protection," *Crop Protection*, 2004, 23:275-285.
- [11] E. C. Oerke, "Crop losses to pests," *Journal of Agricultural Science*, 2006, 144:31-43.
- [12] E. C. Snell-Rood, & D. R. Papaj. "Patterns of phenotypic plasticity in common and rare environments: A study of host use and color learning in the Cabbage White butterfly *Pieris rapae*," *The American Naturalist*, 2009, 173, 615-631.
- [13] FAO, "Global food losses and food waste: extent, causes and prevention," 2011. Available via <http://www.fao.org>. Accessed 12 November 2020.
- [14] FAO/WHO, "The international code of conduct on pesticide management," 2014. Available via http://www.fao.org/fileadmin/templates/agphome/documents/Pests_Pesticides/Code/CODE_2014Sep_ENG.pdf. Accessed 12 November 2020.
- [15] FAO, FAO launches "2020 as the UN's International Year of Plant Health," 2019. Available via <http://www.fao.org/news/story/en/item/1253551/icode/>. Accessed 12 November 2020.
- [16] G. S. Dhaliwal and R. Arora, "An estimate of yield losses due to insect pests in Indian agriculture," *Indian Journal of Ecology*, 1996, 23:70-73.
- [17] G. S. Dhaliwal and R. Arora, "Estimation of losses due to insect pests in field crops," In: Sarath B, Babu KS, Varaprasad K, Anitha RD, Rao VJP, Chakrabarty SK, Chandukar PS (eds) *Resources management in plant protection*. Plant Protection Association of India, Hyderabad, 2002, 1: 11-23.
- [18] G. S. Dhaliwal, R. Arora, and A. K. Dhawan, "Crop losses due to insect pests and determination of economic threshold levels," In: Singh A, Trivedi TP, Sardana HR, Sharma OP, Sabir N (eds) *Recent advances in integrated pest management*. National Centre for Integrated Pest Management, New Delhi, 2003, 12-20pp.
- [19] G. S. Dhaliwal, R. Arora, and A. K. Dhawan, "Crop losses due to insect pests in Indian agriculture: An update," *Indian Journal of Ecology*, 2004, 31:1-7.
- [20] G. S. Dhaliwal, A. K. Dhawan, and R. Singh, "Biodiversity and ecological agriculture: Issues and perspectives," *Indian Journal of Ecology*, 2007, 34:100-109.
- [21] G. S. Dhaliwal, V. Jindal, and A. K. Dhawan, "Insect pest problems and crop losses: Changing trends," *Indian Journal of Ecology*, 2010, 37 (1): 1-7.
- [22] H. H. Cramer, "Plant protection and world crop production," *Bayer Pflanzenschutz-Nachr*, 1967, 20: 1-24.
- [23] K. S. Rao, R. K. Maikhuri, and K. G. Saxena, "Indigenous pest management in the Himalaya," *Soil Biodiversity: Inventory, Functions and Management*, 2015, 29: 383-399.
- [24] K. Tanaka, "Movement of the spiders in arable land," *Plant Protection*, 1989, 43, 1: 34-39.
- [25] K. Rao and B. H. Murthy, *Entomological Society of India [Corporate Author] (eds) "Proceedings of national seminar on crop losses due to insect pests, Hyderabad," Indian Journal of Entomology Vol. I-II (Spl issue), 1983.*
- [26] M. Chandola, S. Rathore, and B. Kumar, "Indigenous pest management practices prevalent among hill farmers of Uttarakhand," *Indian journal of Traditional knowledge*, 2011, 10(2): 311-315.
- [27] M. V. Sampaio, V. H. P. Bueno, L. C. P. Silveira, and A. M. Auad, "Biological Control of Insect Pests in the Tropics," In: Claro KD, Oliveira PS, Gray VR (ed) *Encyclopaedia of*

- Life Support Systems (EOLSS). Developed under the Auspices of the UNESCO, Eolss Publishers, Paris, France. Tropical Biology and Conservation Management, 2009, 3: 29-38. Available via <https://www.eolss.net>. Accessed 12 November 2020.
- [28] O. P. Lal, "Recent advances in Indian entomology," APC, New Delhi, 1996.
- [29] Peters, T. M. "Insects and Human Society," Belchertown, MA: T. M. Peters, 393 Bay Road, 1993.
- [30] R. K. Maikhuri, L. S. Rawat, R. L. Semwal, K. S. Rao, and K. G. Saxena, "Organic farming in Uttarakhand Himalaya, India," International Journal of Ecology and Environmental Sciences, 2015, 41 (3-4): 161-176.
- [31] R. Van den Bosch, P. S. Messenger, and A. P. Gutierrez, "An Introduction to Biological Control," Plenum Press, New York, USA, 1982, pp. 247.
- [32] S. Jayaraj, "Biopesticides and integrated pest management for sustainable crop production," In: Roy NK (ed) Agrochemicals and sustainable agriculture. APC, New Delhi, 1993, pp. 65–81.
- [33] S. Pradhan, "Assessment of losses caused by insect pests of crops and estimation of insect population," In: Pant NC (ed.) Entomology in India, Entomological Society of India, New Delhi, 1964, 17-58pp.
- [34] S. S. Patole, "Review on Beetles (Coleoptera): An agricultural major crop pests of the world," International Journal of Life Sciences, Scienti. Res., 2017, 3 (6): 1424-1432.
- [35] T. Matthias, A. Matthias, and B. Cédric, "Perennial, species-rich wildflower strips enhance pest control and crop yield," Agriculture, Ecosystems and Environment, 2016, 220: 97–103.
- [36] U. Yadav and R. Tiwari, "Effect of smoke on insect mortality and quality parameters of stored wheat at Pantnagar, Uttarakhand," Journal of Entomology and Zoology Studies, 2018, 6 (3): 1661-1666.

Table 1: Checklist of major Insect pests of standing crops

| Order | Family (Number of species) | Insect as a pest (Scientific name) | Host plant | Parts Damaged |
|------------|----------------------------|---|-----------------------------------|----------------------------------|
| Coleoptera | Brentidae (01) | Straight snouted weevil (<i>Trichapion sp.</i>) | Cereals, Vegetables | Root, flower, stem |
| | Chrysomelidae (05) | Cereal Leaf Beetle (<i>Oulema sp</i>) | Cereals | Skeletonize Leaves |
| | | Flea beetle (<i>Altica himalayensis</i>) | Mustard, Lady's finger | Leaf, stems, petals |
| | | Milkweed leaf beetle (<i>Labidomera clivicollis</i>) | Okra, Beans | Leaf, young foliage |
| | | Red Pumpkin beetle (<i>Aulacophora foveicollis</i>) | Cucurbitaceae | Leaf, fruit, flower |
| | | Spotted Cucumber beetle (<i>Diabrotica undecimpunctata</i>) | Cucurbitaceae | Roots, stem, leaf |
| | Curculionidae (01) | Knobbed Weevil (<i>Hadramphus tuberculatus</i>) | Vegetables, Cereals | Root, root hairs, fruit |
| | Elateridae (02) | Black Click beetle (<i>Agriotes gallicus</i>) | Wheat, Potato | Roots, tender stems |
| | | Common Click beetle (<i>Agriotes sputator</i>) | Wheat, Potato | Roots, stems |
| | Meloidae (01) | Blister beetle (<i>Hycleus sp.</i>) | Beans, Tomato, Chilli, Amaranthus | Flower, leaf |
| | Nitidulidae (01) | 4 Spotted Sap beetle (<i>Glischrochilus quadrisignatus</i>) | Corn, Peaches, Tomatoes | Fruits, young foliage |
| | Scarabaeidae (03) | Green June Beetle (<i>Cotinis nitida</i>) | Potato, Onion, | Young foliage of shrubs, flowers |

| | | | | |
|--------------|---------------------|---|--------------------------------|----------------------------------|
| | | May Beetle (<i>Phyllophaga sp.</i>) | Potatoes, Onion, Cucurbitaceae | Young foliage, shrub, flowers |
| | | White Grubs (<i>Holotrichia sp.</i>) | Cole crops, Potato | Roots, tubers |
| Dermaptera | Forficulidae (01) | Earwigs (<i>Forficula sp.</i>) | Cole crops, Cucumber | Foliage, flowers |
| Diptera | Psychodidae (01) | Drain fly (<i>Psychoda alternata</i>) | Beefsteak | Leaf, inflorescence |
| | Tephritidae (01) | Melon fly (<i>Bactrocera cucurbitae</i>) | Bean, Tomato, Brinjal | Fruits, Pods |
| | Tipulidae (01) | Crane-fly (<i>Nephrotoma appendiculata</i>) | Beefsteak, Wheat, | Roots, crown, Leaf |
| Hemiptera | Aphididae (01) | Aphids (<i>Aphis sp.</i>) | Peach, Tomato, Beans, Pulses | Tender buds, stem flower |
| | Coreidae (02) | Coreid bug (<i>Dalader sp.</i>) | Malta, Squash, Pumpkin | Sap, young foliage |
| | | Leaf footed bug (<i>Leptoglossus sp.</i>) | Cucurbits | Sap |
| | Flatidae (01) | Planthopper (<i>Siphanta acuta</i>) | Malta, Eggplant | Leaf, stem, |
| | Miridae (01) | Tarnished plant bug (<i>Lygus sp.</i>) | Beans, Soyabean | Flowers, fruit, leaves |
| | Pentatomidae (01) | Stink/Shield bugs (<i>Nezara viridulag</i>) | Peach, Tomato, Lima beans | Leaf, fruits |
| | Pseudococcidae (01) | Mealybugs (<i>Clavicornus sp.</i>) | Citrus plants, Mulberry | Leaf, buds, tender stems |
| Lepidoptera | Erebidae (01) | Tussock moth (<i>Orgyia sp.</i>) | Amaranth, Maize | Tender stem, leaf |
| | Noctuidae (04) | Cabbage White (<i>Pieris brassicae</i>) | Cabbage, Cauliflower | Defoliation, leaf, |
| | | Cabbage Semilooper (<i>Trichoplusia ni</i>) | Cole crops | Leaf scrabbling, Defoliation |
| | | Fall armyworm (<i>Spodoptera frugiperda</i>) | Maize, Rice | Leaf, whole plant except roots |
| | | Greasy Cutworm (<i>Agrotis ipsilon</i>) | Potato, Tomato, Maize | Leaf, Stem, Tubers |
| | Papilionidae (01) | Citrus Swallowtail (<i>Papilio demodocus</i>) | Sweet Orange | Leaves, young foliage |
| | Sphingidae (01) | Hornworm Caterpillar (<i>Manduca quinquemaculata</i>) | Potato, Tomato, Eggplant | Leaves, flower |
| Orthoptera | Acrididae (02) | Locust (<i>Schistocerca sp.</i>) | Rice, Maize, Vegetables | Leaves, shoot, fruit, stem, bark |
| | | Grasshopper (<i>Cyrtacanthacris tatarica</i>) | Cereals, vegetables | Young leaves |
| Thysanoptera | Thripidae (01) | Thrips (<i>Thrips tabaci</i>) | Onion, Chilli, Cucumber | Leaf, fruits |

Plate 1: Major Insects as a pest in the study site



1



2



3



4



5



6



7



8



9



10



11



12

1. *Pieris brassicae* larvae on *Brassica campestris* 2. *Cotinis nitida* on *Solanum tuberosum* 3. *Holotrichia* sp. 4. Leaves of *Citrus sinensis* damaged by larvae of *Papilio demodocus* 5. *Labidomera clivicollis* 6. *Altica himalayensis* 7. *Nephrotoma appendiculata* on *Triticum aestivum* 8. *Aphis* sp. on *Phaseolus vulgaris* 9. *Orgyia* sp on *Amaranthus* sp. 10. *Bibio* sp. on *Perilla frutescens* 11. Damaged leaves of *Brassica oleracea* 12. *Siphanta acuta*

Plate 2: Natural Enemies of pests in the study site



1



2




3



4



5



6



7



8



9

1. *Coccinellidae* sp. 2. *Halzia* sp. 3. *Coccinella septempunctata* 4. *Odontocolon* sp. (*Ichneumonids*) 5. *Cryptus* sp. 6. *Braconid* sp. 7. *Thomisus* sp. 8. *Mantis* sp. 9. *Mantis religiosa*